



Perpetration of gross human rights violations in South Africa: Association with psychiatric disorders

Dan J Stein, Stacey L Williams, Pamela B Jackson, Soraya Seedat, Landon Myer, Allen Herman, David R Williams

Background. A nationally representative study of psychiatric disorders in South Africa provided an opportunity to study the association between perpetration of human rights violations (HRVs) during apartheid and psychiatric disorder. Prior work has suggested an association between perpetration and post-traumatic stress disorder (PTSD), but this remains controversial.

Methods. Subjects reported on their perpetration of human rights violations, purposeful injury, accidental injury and domestic violence. Lifetime and 12-month prevalence of DSM-IV (*Diagnostic and Statistical Manual*, 4th edition) disorders were assessed with Version 3.0 of the World Health Organization Composite International Diagnostic Interview (CIDI 3.0). Socio-demographic characteristics of these groups were calculated. Odds ratios for the association between the major categories of psychiatric disorders and perpetration were assessed.

Results. HRV perpetrators were more likely to be male, black and more educated, while perpetrators of domestic

violence (DV) were more likely to be female, older, married, less educated and with lower income. HRV perpetration was associated with lifetime and 12-month anxiety and substance use disorders, particularly PTSD. Purposeful and DV perpetration were associated with lifetime and 12-month history of all categories of disorders, whereas accidental perpetration was associated most strongly with mood disorders.

Conclusion. Socio-demographic profiles of perpetrators of HRV and DV in South Africa differ. While the causal relationship between perpetration and psychiatric disorders deserves further study, it is possible that some HRV and DV perpetrators were themselves once victims. The association between accidental perpetration and mood disorder also deserves further attention.

S Afr Med J 2009; 99: 390-395.

South Africa provides a unique laboratory for studying the perpetration of different forms of violence. During apartheid a range of acts of racial violence were perpetrated, including

Department of Psychiatry and Mental Health, University of Cape Town

Dan J Stein, FRCPC, PhD, DPhil

Department of Psychology, East Tennessee State University, Johnson City, Tenn., USA

Stacey L Williams, PhD

Department of Sociology, Indiana University, Bloomington, Ind., USA

Pamela B Jackson, PhD

MRC Stress and Anxiety Disorders Unit, Department of Psychiatry, Stellenbosch University, Tygerberg, W Cape

Soraya Seedat, MB ChB, FCPsych (SA), PhD

School of Public Health and Family Medicine, University of Cape Town

Landon Myer, MB ChB, PhD

National School of Public Health, University of Limpopo (Medunsa Campus), Pretoria

Allen Herman, MB ChB, PhD

Department of Society, Human Development and Health, Harvard School of Public Health, and Department of African and African American Studies, Harvard University, Cambridge, Mass., USA

David R Williams, MPH, PhD

gross human rights violations (HRVs). However, both before and after apartheid there have also been significant levels of domestic and criminal violence.¹ Injuries were the second-leading cause of premature death and interpersonal injury and accounted for 14.3% of disability-adjusted life-years in South Africa in 2000,² with interpersonal injury predominating.³ A survey in 1998 found that 2 in 1 000 South Africans had experienced a violent injury requiring medical treatment in the previous month, while 9 in 1 000 households had had a violent death in the previous year.⁴ Community surveys have emphasised high levels of domestic and partner violence,⁵⁻¹¹ including sexual assault and rape.¹²⁻¹⁶

Studies have suggested significant associations between perpetration of violence and mental illness. Although public perceptions of the link between disorders such as schizophrenia and violence are clouded by stigma, a systematic review found perpetration of violence and violence victimisation to be more common in persons with severe mental illness than in the general population.¹⁷ The relationship among substance use, violence and victimisation is also well established, particularly the association between men's substance use and perpetration of physical violence.^{18,19} While associations between perpetration and antisocial traits²⁰⁻²⁴ can be expected, there is also the less obvious possibility that many perpetrators have themselves been exposed to trauma (e.g. childhood abuse, marital violence, military combat), or suffer from post-traumatic stress disorder (PTSD).^{16,24-30}

Corresponding author: D J Stein (dan.stein@uct.ac.za)



The South African Stress and Health (SASH) study is a nationally representative survey that provides information on the prevalence of common mental disorders and on a range of other variables, including perpetration of violence.³¹ We report on the association between perpetration and common mental disorders.

Methods

Subject sample

The SASH was a national probability sample of adult South Africans living in households and hostel quarters, obtained between January 2002 and June 2004.³¹ Hostel quarters were included to maximise coverage of young working-age males, but did not include individuals in institutions or the military. The sample was selected using a three-stage probability sample design. A total sample of 5 089 households was selected for the SASH; field interviews were obtained with 4 433 (87.1%) and based on quality control criteria, 4 351 were retained for analysis.

Diagnostic interview

The diagnostic interview used in the SASH was the World Health Organization (WHO) Composite International Diagnostic Interview Version 3.0 (CIDI 3.0),³² a fully structured lay-administered interview that generates diagnoses according to the criteria of both the ICD-10 (*International Statistical Classification of Diseases and Related Health Problems*, 10th revision) and DSM-IV (*Diagnostic and Statistical Manual*, 4th edition) diagnostic systems. Because of time constraints, the interview excluded several disorders (e.g. specific phobia, impulse control disorders other than intermittent explosive disorder). DSM-IV criteria are used in this report. Given the potential significance of PTSD, the analyses focus on this specific disorder and four summary categories of mental disorders: anxiety disorders (panic disorder, agoraphobia, social phobia, generalised anxiety disorder, PTSD), mood disorders (major depressive disorder, dysthymia), substance use disorders (alcohol abuse, alcohol dependence, drug abuse, drug dependence), and a global category of any of the abovementioned disorders. DSM-IV organic exclusion rules and diagnostic hierarchy rules were applied to all diagnoses, except in the case of substance use disorders where abuse was defined with or without dependence. Interviewers were trained in the administration of the CIDI in centralised group sessions lasting 1 week. The interviews were conducted face to face in English, Afrikaans, Zulu, Xhosa, Northern Sotho, Southern Sotho or Tswana. The protocol was approved by an ethics committee and all subjects gave informed consent. Interviews lasted an average of 3½ hours, with some requiring more than one visit to complete.

Assessment of perpetration

Perpetration was assessed using different probes. First, respondents were asked: 'Because of political reasons did you ever arrest or detain someone, kidnap or abduct someone, participate in the destruction of someone's home or property, physically beat or injure someone, participate in the death of someone, kill someone, participate in any other form of torture or human rights violation?' Our HRV perpetrator variable is a dichotomous indicator contrasting individuals who reported one or more of those experiences to those who reported none (HRV perpetration). Second, in screening for PTSD, subjects were asked whether they had ever seriously injured, tortured or killed someone (purposeful perpetration), or accidentally seriously injured or killed someone (accidental perpetration). Third, domestic violence perpetration was assessed by the frequency with which the respondent had slapped or hit, thrown something at, or pushed, grabbed or shoved her/his current or former spouse or partner.³³

Statistical analysis

The person-level SASH data were weighted to adjust for differential probabilities of selection within households, differential non-response, and residual discrepancies between the sample and the population on a profile of census demographic and geographical variables and used in all data analyses. Data analysis was carried out using SAS and SAS-callable SUDAAN software to adjust estimates of statistical significance for the weighting and clustering of the data.

For each perpetrator variable, bivariate comparisons of socio-demographic variables (gender, age, race, income, marital status, education, employment status, location) were undertaken, and then associations of perpetration with mental disorder were assessed, adjusting for these socio-demographic variables. (Use of race variables is not intended to reify socially constructed categories, but to allow an exploration of the effects of historical circumstances on current public health issues.)

Results

Domestic violence (DV) was common (15%), with lower figures for HRV perpetration (3%), and purposeful (1.0%) and accidental (1.9%) perpetration. Table I demonstrates the bivariate comparisons with socio-demographic factors. HRV perpetrators were more likely to be male, black, and better educated. Purposeful perpetrators were more likely to be white and employed with a trend towards being male, while accidental perpetrators were more likely to be male. DV perpetrators were more likely to be female, older, married, less educated, and with lower income.

Logistic regression models are presented for each of the perpetration variables. Table II presents the association between HRV perpetration and mental disorders. As in the subsequent tables, each row of the table presents findings from a



Table I. Demography of perpetration

	Perpetrator of HRV	Purposeful perpetrator of injury	Accidental perpetrator of injury	Perpetrator of domestic violence
Total				
N (%)	111 (3.0)	38 (1.0)	75 (1.9)	690 (15.7)
Gender				
Male	78 (4.8)	24 (1.4)	43 (2.6)	242 (14.1)
Female	33 (1.4)	14 (0.5)	32 (1.3)	448 (17.1)
$\chi^2 (p)$	16.5 (0.000)	3.6 (0.064)	11.0 (0.002)	4.0 (0.050)
Race				
Black	97 (3.3)	28 (0.9)	63 (2.1)	507 (14.8)
Coloured	9 (1.8)	5 (1.1)	6 (1.4)	105 (19.5)
White	4 (2.6)	5 (1.4)	3 (0.8)	42 (15.8)
Indian	1 (0.5)	0 (0.0)	3 (1.8)	36 (24.7)
$\chi^2 (p)$	16.1 (0.002)	25.5 (0.000)	4.3 (0.245)	7.7 (0.064)
Age (yrs)				
18 - 34	53 (2.6)	22 (1.0)	32 (1.9)	224 (10.1)
35 - 49	44 (4.1)	11 (1.3)	30 (2.3)	276 (20.8)
50 - 64	10 (2.5)	4 (0.5)	12 (1.6)	139 (23.0)
≥65	4 (2.2)	1 (0.3)	1 (0.6)	50 (24.1)
$\chi^2 (p)$	3.6 (0.322)	8.3 (0.049)	5.5 (0.150)	63.7 (0.000)
Marital status				
Unmarried	56 (2.8)	24 (1.3)	37 (1.8)	143 (5.7)
Married	55 (3.2)	14 (0.7)	38 (2.0)	547 (26.0)
$\chi^2 (p)$	0.4 (0.519)	1.8 (0.189)	0.1 (0.748)	165 (0.000)
Education				
None	8 (2.8)	5 (0.9)	7 (2.2)	68 (24.7)
Grade 1 - 7	18 (2.4)	5 (0.6)	19 (2.3)	196 (22.1)
Grade 8 - 11	41 (3.1)	12 (1.0)	27 (1.8)	256 (15.5)
Grade 12	20 (1.9)	11 (1.4)	10 (1.4)	92 (11.1)
Grade 13+	24 (4.8)	5 (0.6)	12 (2.1)	78 (11.8)
$\chi^2 (p)$	11.9 (0.026)	3.2 (0.523)	1.4 (0.836)	49.6 (0.000)
Income (R)				
0	14 (2.9)	6 (0.7)	11 (1.8)	102 (15.9)
1 - 1 500	28 (2.7)	8 (0.9)	14 (1.2)	178 (17.9)
1 501 - 16 500	21 (2.7)	11 (1.3)	13 (1.4)	156 (17.6)
16 501 - 97 500	20 (2.6)	5 (0.8)	16 (1.7)	144 (15.3)
≥97 501	28 (4.0)	8 (1.0)	21 (3.2)	110 (12.1)
$\chi^2 (p)$	2.4 (0.661)	1.6 (0.800)	7.9 (0.110)	13.2 (0.016)
Employment status				
Unemployed	72 (2.8)	20 (0.6)	52 (1.7)	449 (13.9)
Employed	39 (3.4)	18 (1.6)	23 (2.3)	241 (19.7)
$\chi^2 (p)$	0.7 (0.405)	4.4 (0.041)	0.8 (0.379)	7.5 (0.008)
Location				
Rural	43 (2.7)	14 (0.6)	33 (2.0)	316 (17.2)
Urban	68 (3.1)	24 (1.1)	42 (1.8)	374 (14.8)
$\chi^2 (p)$	0.4 (0.510)	3.6 (0.063)	0.2 (0.679)	2.2 (0.146)

multivariate regression model in which the association between the perpetrator variable and disorder was examined adjusted for all of the socio-demographic factors. HRV perpetration was associated with lifetime and 12-month anxiety and substance use disorders. The odds ratios (ORs) were particularly large for PTSD. Purposeful perpetration (Table III) was associated with lifetime and 12-month history of all categories of disorders. Despite the small numbers, the ORs were very large for both 12-month and lifetime rates of disorder, suggesting a robust association. In contrast, the association with disorders is much

weaker for accidental perpetration (Table IV), although there is an elevated risk for lifetime and 12-month mood disorder and 12-month anxiety and substance disorders. DV perpetration (Table V) was associated with lifetime and 12-month history of all categories of disorders.

Discussion

We found that different kinds of perpetration appear to be associated with different socio-demographic profiles, and are associated with different clinical disorders. There have



Table II. Multivariate odds ratios (with 95% confidence intervals) for DSM-IV disorders (predictor: perpetration of human rights violations, model N=4 188)

Outcome	OR	LCI	UCI	<i>p</i>
Lifetime disorders				
All DSM-IV	3.12	1.89	5.12	0.000
Anxiety disorder	2.80	1.69	4.65	0.000
PTSD	5.82	2.67	12.72	0.000
Mood disorder	1.44	0.67	3.12	0.345
Substance use	3.18	2.11	4.81	0.000
Past 12-month disorders				
All DSM-IV	1.52	0.95	2.44	0.082
Anxiety disorder	1.88	0.98	3.61	0.058
PTSD	5.24	1.68	16.35	0.005
Mood disorder	0.50	0.16	1.57	0.232
Substance use	1.97	1.01	3.83	0.048

LCI = lower confidence interval; UCI = upper confidence interval.

Table III. Multivariate odds ratios (with 95% confidence intervals) for DSM-IV disorders (predictor: purposeful perpetration of serious injury, torture or death, model N=4 310)

Outcome	OR	LCI	UCI	<i>p</i>
Lifetime disorders				
All DSM-IV	6.72	3.07	14.68	0.000
Anxiety disorder	5.63	2.84	11.17	0.000
PTSD	5.19	1.64	16.37	0.006
Mood disorder	4.88	1.71	13.88	0.004
Substance use	2.91	1.20	7.04	0.019
Past 12-month disorders				
All DSM-IV	6.09	2.53	14.63	0.000
Anxiety disorder	7.38	2.14	25.50	0.002
PTSD	22.92	6.62	79.40	0.000
Mood disorder	4.49	1.29	15.59	0.019
Substance use	2.47	1.09	5.60	0.031

LCI = lower confidence interval; UCI = upper confidence interval.

been few population-based surveys of the clinical correlates of perpetration, and these findings have implications for understanding the nature of perpetration, and for developing appropriate interventions.

HRV perpetrators were more likely to be male, black and more educated, and HRV perpetration was associated with lifetime and 12-month anxiety and substance use disorders, particularly PTSD. These subjects may have been employed by the security forces (police, military, etc.) during apartheid. While causal relationships cannot be determined from these data, studies have indicated that perpetration of violent acts during combat can lead to subsequent PTSD.³⁴

In contrast, DV perpetrators were more likely to be female, older, married, less educated, and with lower income. Although statistical significance was not reached, there was a trend for race to be significant with elevated risk associated with being Indian or coloured. DV perpetration was also associated with lifetime

and 12-month history of all categories of disorders. The gender findings are consistent with previous studies; in South Africa, partner violence in adolescents was higher in females than in males,⁷ and a meta-analysis of over 80 studies found that women were more likely than men to perpetrate violent behaviour in their intimate relationships than men.³⁵ However, it is important to emphasise that when gender analyses include differences in impact and context, women are disproportionately victimised by partner violence.^{36,37} Men are more coercively controlling of their partners than females – a context which may significantly change the meaning of females' use of violence in a relationship.³⁸ Similar considerations may also apply to other socio-demographic correlates of DV perpetration found here. In keeping with the association of DV perpetration with mental disorders, studies have found a range of psychopathology in this population.^{21-24,38-42}

There were small samples of purposeful and accidental perpetrators. Purposeful perpetrators were more likely to be white, under 50 and employed, and may represent those employed by the security establishment and involved in particularly serious acts of perpetration. In contrast, apart from being male, accidental perpetrators had no significant socio-demographic associations. Purposeful perpetration was associated with lifetime and 12-month history of all categories of disorders, whereas accidental perpetration was associated most strongly with mood disorders. These findings are consistent with the high levels of psychopathology in members of the South African security forces,⁴³ and with descriptions of regret and remorse in accidental perpetrators.

A number of limitations must be emphasised. First, there are concerns about whether self-report questions about perpetration are answered truthfully. Partial reassurance is perhaps obtained from retrospective reporting of sensitive questions such as that of childhood adversity; although false-negative reports are common, false-positive reports are rare, so that retrospective case-control studies of such adversities are potentially valid.⁴⁴ We also implemented procedures known to improve the accuracy of reporting of sensitive questions in surveys.⁴⁵ These included using a commitment probe (having respondents pledge to trying hard to answer the questions accurately) and using 'forgiving' language (the questions about human rights perpetration were preceded by a statement that said, 'during times of conflict and because of one's job, people sometimes do things that they normally would not do'). Nevertheless, while many respondents admitted to DV perpetration, consistent with previous work in South Africa,⁵⁻¹⁰ few admitted to purposeful or accidental perpetration, and these findings should therefore be interpreted with caution. Second, these analyses did not adjust for levels of severity of perpetration; more severe and less severe perpetration may have different socio-demographic and clinical correlates.^{36,46} Other aspects of perpetration were also not captured (for

**Table IV. Multivariate odds ratios (with 95% confidence intervals) for DSM-IV disorders (predictor: accidental perpetration of serious injury or death, model N=4 293)**

Outcome	OR	LCI	UCI	p
Lifetime disorders				
All DSM-IV	1.99	1.17	3.37	0.012
Anxiety disorder	1.36	0.70	2.65	0.354
PTSD	0.42	0.05	3.29	0.399
Mood disorder	2.45	1.31	4.58	0.006
Substance use	1.75	0.92	3.33	0.088
Past 12-month disorders				
All DSM-IV	2.66	1.42	4.98	0.003
Anxiety disorder	2.26	1.03	4.93	0.041
PTSD	-	-	-	-
Mood disorder	3.72	1.57	8.79	0.003
Substance use	2.16	1.00	4.69	0.051

LCI = lower confidence interval; UCI = upper confidence interval.

Table V. Multivariate odds ratios (with 95% confidence intervals) for DSM-IV disorders (predictor: perpetration of domestic violence, model N=4 136)

Outcome	OR	LCI	UCI	p
Lifetime disorders				
All DSM-IV	2.00	1.67	2.39	0.000
Anxiety disorder	1.47	1.09	1.96	0.011
PTSD	2.02	1.22	3.36	0.007
Mood disorder	1.84	1.34	2.54	0.000
Substance use	2.43	1.76	3.36	0.000
Past 12-month disorders				
All DSM-IV	1.80	1.40	2.31	0.000
Anxiety disorder	1.58	1.12	2.25	0.011
PTSD	2.73	0.92	8.05	0.069
Mood disorder	1.76	1.11	2.78	0.017
Substance use	2.09	1.39	3.14	0.001

LCI = lower confidence interval; UCI = upper confidence interval.

example, the extent to which DV perpetration was one-sided or mutual³⁸). Third, associations between perpetration and mental disorder were not investigated separately in males and females; previous work has suggested gender-specific correlations.^{21,22,41} Fourth, the data do not allow temporal or causal relationships to be established, and the extent to which mental disorders are causes or consequences of perpetration cannot be determined.

Nevertheless, this study provides the first data on associations between multiple types of perpetration and mental illness in South Africa, and provides avenues for exploration by researchers and policy makers, locally and abroad.

Several interventions are effective for reducing interpersonal violence.⁴⁷⁻⁵⁰ Given the high burden of disease associated with trauma in general in South Africa, and interpersonal violence in particular, and the relationships of interpersonal violence with mental disorder and sexually transmitted infections such as HIV/AIDS,^{13,51,52} these interventions deserve serious consideration. The limitations of the data here need to be taken

into account when formulating such interventions; for example, the more coercive perpetration of males probably deserves priority. The high prevalence of domestic violence perpetration warrants special attention, but an understanding of the nature of other kinds of perpetration and the development of appropriate interventions is also key.^{53,54}

Acknowledgements. The South African Stress and Health study was carried out in conjunction with the World Health Organization World Mental Health (WMH) Survey Initiative. We thank the WMH staff for assistance with instrumentation, fieldwork, and data analysis. These activities were supported by the US National Institute of Mental Health (R01MH070884), the John D and Catherine T MacArthur Foundation, the Pfizer Foundation, the US Public Health Service (R13-MH066849, R01-MH069864, and R01 DA016558), the Fogarty International Center (FIRCA R01-TW006481), the Pan American Health Organization, Eli Lilly and Company, Ortho-McNeil Pharmaceutical, Inc., GlaxoSmithKline, and Bristol-Myers Squibb. The South Africa Stress and Health study was funded by grant R01-MH059575 from the National Institute of Mental Health and the National Institute of Drug Abuse with supplemental funding from the South African Department of Health and the University of Michigan. Dan Stein and Soraya Seedat are also supported by the Medical Research Council of South Africa. A complete list of WMH publications can be found at <http://www.hcp.med.harvard.edu/wmh/>

The authors thank Kathleen McGaffigan (Harvard University) for her assistance with the statistical analyses.

References

- Stein DJ, Seedat S, Emsley RA. Violence in the world and in South Africa – what is a doctor to do? *S Afr Med J* 2002; 92: 789-790.
- Norman R, Matzopoulos R, Groenewald P, Bradshaw D. The high burden of injuries in South Africa. *Bull World Health Organ* 2007; 85: 695-702.
- Matzopoulos R, Seedat M. National Injury Mortality Surveillance System – a call for city-level injury prevention. *S Afr Med J* 2005; 95: 708.
- Doolan K, Ehrlich R, Myer L. Experience of violence and socioeconomic position in South Africa: a national study. *Plos One* 2007; 2: e1290.
- Jewkes R, Levin J, Penn-Kekana L. Risk factors for domestic violence: findings from a South African cross-sectional study. *Soc Sci Med* 2002; 55: 1603-1617.
- Abrahams N, Jewkes R, Laubscher R, Hoffman M. Intimate partner violence: prevalence and risk factors for men in Cape Town, South Africa. *Violence Vict* 2006; 21: 247-264.
- Flisher AJ, Myer L, Merais A, Lombard C, Reddy P. Prevalence and correlates of partner violence among South African adolescents. *J Child Psychol Psychiatry* 2007; 48: 619-624.
- Dinan BA, McCall GJ, Gibson D. Community violence and PTSD in selected South African townships. *J Interpers Violence* 2004; 19: 727-742.
- Dunkle KL, Jewkes RK, Nduna M, et al. Perpetration of partner violence and HIV risk behaviour among young men in the rural Eastern Cape, South Africa. *AIDS* 2006; 20: 2107-2114.
- Dunkle KL, Jewkes RR, Brown HC, et al. Gender-based violence, relationship power, and risk of HIV infection in women attending antenatal clinics in South Africa. *Lancet* 2004; 363: 1415-1421.
- Swart LA, Stevens MSG, Ricardo I. Violence in adolescents' romantic relationships: findings from a survey amongst school-going youth in a South African community. *J Adolesc* 2002; 25: 385-395.
- Jewkes R, Dunkle K, Koss MP, et al. Rape perpetration by young, rural South African men: Prevalence, patterns and risk factors. *Soc Sci Med* 2006; 63: 2949-2961.
- Kalichman SC, Simbayi LC, Cain D, Cherry C, Henda N, Cloete A. Sexual assault, sexual risks and gender attitudes in a community sample of South African men. *AIDS Care* 2007; 19: 20-27.
- Jewkes R, Abrahams N. The epidemiology of rape and sexual coercion in South Africa: an overview. *Soc Sci Med* 2002; 55: 1231-1244.
- Buga GAB, Amoko DHA, Ncayiyana DJ. Sexual behaviour, contraceptive practice and reproductive health among school adolescents in rural Transkei. *S Afr Med J* 1996; 86: 523-527.
- Andersson N, Ho-Foster A, Matthis J, et al. National cross sectional study of views on sexual violence and risk of HIV infection and AIDS among South African school pupils. *BMJ* 2004; 329: 952-954A.



17. Choe JY, Teplin LA, Abram KM. Perpetration of violence, violent victimization, and severe mental illness: Balancing public health concerns. *Psychiatr Serv* 2008; 59: 153-164.
18. Testa M. The role of substance use in male-to-female physical and sexual violence – a brief review and recommendations for future research. *J Interpers Violence* 2004; 19:1494-1505.
19. Cunradi CB, Caetano R, Schafer J. Alcohol-related problems, drug use, and male intimate partner violence severity among US couples. *Alcohol Clin Exp Res* 2002; 26: 493-500.
20. Capaldi DM, Owen LD. Physical aggression in a community sample of at-risk young couples: Gender comparisons for high frequency, injury, and fear. *J Fam Psychol* 2001; 15: 425-440.
21. Yang M, Coid J. Gender differences in psychiatric morbidity and violent behaviour among a household population in Great Britain. *Soc Psychiatry Psychiatr Epidemiol* 2007; 42: 599-605.
22. Magdol L, Moffitt TE, Caspi A, Newman DL, Fagan J, Silva PA. Gender differences in partner violence in a birth cohort of 21-year-olds: Bridging the gap between clinical and epidemiological approaches. *J Consult Clin Psychol* 1997; 65: 68-78.
23. Feder L, Henning K. A comparison of male and female dually arrested domestic violence offenders. *Violence Vict* 2005; 20: 153-171.
24. Stuart GL, Moore TM, Gordon KC, Ramsey SE, Kahler CW. Psychopathology in women arrested for domestic violence. *J Interpers Violence* 2006; 21: 376-389.
25. Taft CT, Pless AP, Stalans LJ, Koenen KC, King LA, King DW. Risk factors for partner violence among a national sample of combat veterans. *J Consult Clin Psychol* 2005; 73: 151-159.
26. Fehon DC, Grilo CM, Lipschitz DS. A comparison of adolescent inpatients with and without a history of violence perpetration – impulsivity, PTSD, and violence risk. *J Nerv Ment Dis* 2005; 193: 405-411.
27. Kim YS, Leventhal BL, Koh YJ, Hubbard A, Boyce WT. School bullying and youth violence – causes or consequences of psychopathologic behavior? *Arch Gen Psychiatry* 2006; 63: 1035-1041.
28. Swan SC, Snow DL. The development of a theory of women's use of violence in intimate relationships. *Violence against Women* 2006; 12: 1026-1045.
29. Abrahams N, Jewkes R. Effects of South African men's having witnessed abuse of their mothers during childhood on their levels of violence in adulthood. *Am J Public Health* 2005; 95: 1811-1816.
30. Abel EM. Comparing the social service utilization, exposure to violence, and trauma symptomatology of domestic violence female 'victims' and female 'batterers'. *J Fam Violence* 2001; 16: 401-420.
31. Williams DR, Herman A, Kessler RC, et al. The South Africa Stress and Health Study: rationale and design. *Metab Brain Dis* 2004; 19: 135-147.
32. Kessler RC, Ustun TB. The World Mental Health (WMH) Survey Initiative Version of the World Health Organization (WHO) Composite International Diagnostic Interview (CIDI). *Int J Methods Psychiatr Res* 2004; 13: 93-121.
33. Straus MA. Measuring intrafamily conflict and violence: The Conflict Tactics (CT) Scales. *Journal of Marriage and the Family* 1979; 41: 75-88.
34. Seedat S, le Roux C, Stein DJ. Prevalence and characteristics of trauma and post-traumatic stress symptoms in operational members of the South African National Defence Force. *Mil Med* 2003; 168: 71-75.
35. Archer J. Sex differences in aggression between heterosexual partners: A meta-analytic review. *Psychol Bull* 2000; 126: 651-680.
36. Hamberger LK. Men's and women's use of intimate partner violence in clinical samples: Towards a gender-sensitive analysis. *Violence Vict* 2005; 20: 131-151.
37. Cogan R, Fennell T. Sexuality and the commission of physical violence to partners and non-partners by men and women. *J Consult Clin Psychol* 2007; 75: 960-967.
38. Johnson MP. Conflict and control: Gender symmetry and asymmetry in domestic violence. *Violence against Women* 2006; 12: 10013-1018.
39. Paterson J, Feehan M, Butler S, Williams M, Cowley-Malcolm ET. Intimate partner violence within a cohort of Pacific mothers living in New Zealand. *J Interpers Violence* 2007; 22: 698-721.
40. Henning K, Jones A, Holdford R. Treatment needs of women arrested for domestic violence – a comparison with male offenders. *J Interpers Violence* 2003; 18: 839-856.
41. Caetano R, Cunradi C. Intimate partner violence and depression among whites, blacks, and Hispanics. *Ann Epidemiol* 2003; 13: 661-665.
42. Dinwiddie SH. Psychiatric disorders among wife batterers. *Compr Psychiatry* 1992; 33: 411-416.
43. Emsley RA, Seedat S, Stein DJ. Posttraumatic stress disorder and occupational disability in South African security force members. *J Nerv Ment Dis* 2003; 191: 237-241.
44. Hardt J, Rutter M. Validity of adult retrospective reports of adverse childhood experiences: review of the evidence. *J Child Psychol Psychiatry* 2004; 45: 260-273.
45. Tourangeau R, Yan T. Sensitive questions in surveys. *Psychol Bull* 2007; 133: 859-883.
46. Williams SL, Frieze IH. Patterns of violent relationships, psychological distress, and marital satisfaction in a national sample of men and women. *Sex Roles* 2005; 52: 771-784.
47. Krug EG, Dahlberg LL, Mercy JA, Zwi A, Lozano R. *World Report on Violence and Health*. Geneva: World Health Organization, 2002.
48. Kim JC, Watts CH, Hargreaves JR, et al. Understanding the impact of a microfinance-based intervention on women's empowerment and the reduction of intimate partner violence in South Africa. *Am J Public Health* 2007; 97: 1794-1802.
49. Pronyk PM, Hargreaves JR, Kim JC, et al. Effect of a structural intervention for the prevention of intimate-partner violence and HIV in rural South Africa: a cluster randomised trial. *Lancet* 2006; 368: 1973-1983.
50. Usdin S, Scheepers E, Goldstein S, Japhet G. Achieving social change on gender-based violence: a report on the impact evaluation of Soul City's fourth series. *Soc Sci Med* 2005; 61: 2434-2445.
51. Jewkes R, Dunkle K, Nduna M, et al. Factors associated with HIV sero-status in young rural South African women: connections between intimate partner violence and HIV. *Int J Epidemiol* 2006; 35: 1455-1460.
52. Andersson G, Carlbring P, Kaldö V, Strom L. Screening of psychiatric disorders via the Internet. A pilot study with tinnitus patients. *Nord J Psychiatry* 2004; 58: 287-291.
53. Nell V. Cruelty's rewards: The gratifications of perpetrators and spectators. *Behav Brain Sci* 2006; 29: 211-224.
54. Kaminer D, Stein DJ. Sadistic personality disorder in perpetrators of human rights abuses: a South African case study. *J Personal Disord* 2001; 15: 475-486.

Accepted 2 February 2009.