



## Unsuitable blood glucose measuring devices in neonatal or paediatric acute care

**To the Editor:** In acute neonatal or paediatric care, bedside blood glucose measurement is aimed predominantly at identifying hypoglycaemia, but high readings prompt the suspicion of diabetes mellitus or the stress hyperglycaemia of critical illness, in which high blood glucose levels correlate positively with poorer outcome.<sup>1</sup>

A recent case has reminded us that bedside glucose testing is not always specific for glucose only.<sup>2</sup>

A 13-day-old girl presented with persistent neonatal jaundice, profound micro-angiopathic haemolytic anaemia, a grossly prolonged prothrombin time and a white cell leukaemoid reaction of  $53 \times 10^9/l$ . *Escherichia coli* was cultured from the urine, and the patient responded to antibiotics.

Blood glucose monitoring with Accu-chek glucose meter (Roche Diagnostics) test strips gave persistently high readings. The absence of glycosuria and a comparison with laboratory blood glucose values alerted us to the fact that the readings were falsely elevated. This was due to a high blood galactose level in this patient, in whom the diagnosis of galactosaemia due to transferase deficiency was subsequently confirmed. We had failed to appreciate the fact that the glucose dye oxidoreductase mediator reaction (glucose dehydrogenase) as used in the Roche Accu-chek also measures galactose.<sup>3</sup>

Previously, in a comparison of point-of-care glucose meters Newman *et al.*<sup>4</sup> had found the Roche Accu-chek to give equimolar interference of galactose with glucose and therefore to risk gross overestimation of glucose in cases of galactosaemia.

A normal blood galactose level ( $<0.44 \text{ mmol/l}^5$ ) is not high enough to result in clinically significant elevation of the combined blood glucose and galactose level. However, permanent or transient elevation of blood galactose is found in galactosaemia, liver dysfunction or maturational delay of galactose transport or utilisation,<sup>5</sup> and then can indeed affect measurements and clinical judgement. The present case illustrates this risk to the detriment of the patient.

We conclude that the Accu-chek glucose meter is not suitable as a blood glucose measuring device in neonatal or paediatric intensive care units wherever elevation of blood galactose levels cannot be excluded *a priori*, and should be removed from routine use in such areas.

**D F Wittenberg**  
**A Terblanche**  
**I Smuts**  
**J Opperman**

Department of Paediatrics and Child Health  
Pretoria Academic Hospital and  
University of Pretoria  
dwittenb@medic.up.ac.za

1. Mariscalco M. Hyperglycemia in children: unique challenges ... unanswered questions. *Pediatr Crit Care Med* 2005; 6(4): 498-499.
2. Jacobsz M, Dednam M. Pre-analytical, analytical and post-analytical considerations in glucose point-of-care testing (letter). *S Afr Med J* 2007; 97: 805.
3. Accu-chek® Active Package Insert. User information issued 2006-03. Roche Diagnostics, Singapore.
4. Newman JD, Ramsden CA, Balazs ND. Monitoring neonatal hypoglycemia with the Accu-chek advantage II glucose meter: the cautionary tale of galactosemia [Letter]. *Clin Chem* 2002; 48(11): 2071.
5. Ono H, Mawatari H, Mizoguchi N, *et al.* Transient galactosemia detected by neonatal mass screening. *Pediatr Internat* 1999; 41: 281-284.

## Information for patients

**To the Editor:** Most patients crave information concerning their illnesses and the interventions planned for their treatment. Most doctors are too busy to be able to give patients the additional time necessary to impart this and to encourage the patient to ask questions. My experience as the National Medical Ombudsman confirmed how important communication was in preventing subsequent problems. Furthermore, the law requires that for consent for a procedure to be valid, the patient must be fully informed – this is hardly ever done.

For all these reasons I would urge the various societies, specialist and general practice, to prepare relevant 'hand-outs'



**Your 1 stop medical mall**



[www.medmedical.co.za](http://www.medmedical.co.za)

All orders placed via our website will receive a  
10% discount on completion

Established 1983 Revived in 2007

With over 3000 products  
we know we can supply you with  
the best products, best service, best advice  
at the right price.

Tuxedo Junction  
4 Dunottar Street  
Orange Grove  
(off Louis Botha)

[info@medmedical.co.za](mailto:info@medmedical.co.za)  
011 640 5262

Come visit our new showroom  
in Tuxedo Junction

1231





for their patients, spelling out what is the matter with them, what, if anything, caused the problem, what is proposed to be done, what the possible complications are (with incidences) and how the post-intervention period is likely to run.

I anticipate that one objection to this suggestion will be that every patient is unique and that generalisations cannot therefore be made; another that every doctor is different and has different approaches to problems. I would counter these objections, firstly by claiming that there are many routine procedures in every branch of medicine and that it is in fact

possible to make broad generalisations, which, if necessary, can be modified for individual patients. Secondly, the societies could put out generalised 'hand-outs' in digital form which could than be modified by the individual practitioners.

#### Olliver Ransome

PO Box 346  
Ramsgate  
4285  
oransome@eject.co.za

# WEB-BASED CPD short courses in PHARMACOLOGICAL Principles of DRUG THERAPY (distance learning) – may lead to an Hons B Sc degree in Pharmacology

#### Enrol throughout the year

This unique Internet and Web-assisted learning programme has been developed by the School of Pharmacy (Faculty of Health Sciences) to give medical practitioners a general insight into the basic pharmacological and ethical principles on which therapeutic strategies are based, enabling them to practise more effective pharmacotherapy in community medicine.

Enrol for the whole programme or selected modules only.

#### Who should enrol?

- Medical Practitioners and Specialists
- Pharmacists

#### Why enrol?

- Continued Professional Development (CPD)
- Stay abreast with the latest trends in Pharmacology
- Empowerment for career development

Candidates can either enrol for the degree programme or specific module(s) as certificate course(s).

#### Some of the certificate courses on offer for you:

- Drugs and the Peripheral Nervous System
- Drugs and the Central Nervous System
- Drugs and the Renal and Cardiovascular Systems
- Drugs for Pain, Inflammation, Fever and Airways Disease
- Chemotherapy of Infections and Infestations
- Hormones, Hormonoids and Hormone Antagonists Drugs for GIT and Skin
- Vitamins, Haematopietics and Immunopharmacology
- Advanced Pharmacological Principles
- Evidence-Based Medicine

#### Please visit our website at:

[www.nwu.ac.za/webpharmacol](http://www.nwu.ac.za/webpharmacol) for more information  
or contact: Alsorika Kraukamp  
Tel: +27(0)18 299 2260 Fax: +27(0)18 299 4244  
E-mail: [alsorika.kraukamp@nwu.ac.za](mailto:alsorika.kraukamp@nwu.ac.za)

BLACK KHAKI 070694

