

an adequate response, but there may be better biomarkers, e.g. soluble transferrin receptors.²

Consequently, there is a need for guidelines that are clearly evidence-based, depend on multiple biomarkers, and take cognisance of varied conditions, and we call for further work and collaboration across disciplines.²

1. Linegar AG, Knottenbelt JD, Wormald PJ. Accuracy of a portable haemoglobinometer in clinical practice. *S Afr Med J* 1991;79: 547-548.
2. Clark SF. Iron deficiency anemia: diagnosis and management. *Curr Opin Gastroenterol* 2009; 25: 122-128.

3. Discombe G, Goddard AJ, Hill ID. The accuracy of haemoglobin determinations with the Spencer haemoglobinometer. *J Coll Gen Pract* 1966; 12: 251-252.
4. Neville RG. Evaluation of portable haemoglobinometer in general practice. *Br Med J (Clin Res Ed)* 1987; 294: 1263-1265.
5. Adamson EA, Bailey GR, Richards N, Wilson H. Prevalence of anaemia in an inner city primary school population. *Arch Dis Child* 2008; 93: 453.
6. Frewin R, Henson A, Provan D. ABC of clinical haematology. Iron deficiency anaemia. *BMJ* 1997; 314: 360-363.
7. Northrop-Clewes CA. Interpreting indicators of iron status during an acute phase response – lessons from malaria and human immunodeficiency virus. *Ann Clin Biochem* 2008; 45: 18-32.
8. Smellie WS, Forth J, Bareford D, et al. Best practice in primary care pathology: review 3. *J Clin Pathol* 2006; 59: 781-789.
9. Vannella L, Aloe Spiriti MA, Cozza G, et al. Benefit of concomitant gastrointestinal and gynaecological evaluation in premenopausal women with iron deficiency anaemia. *Aliment Pharmacol Ther* 2008; 28: 422-430.

Errata

In the article by A J Okwi *et al.* which appeared on pp. 887 - 891 of the December 2009 *SAMJ*, the first part of the title should have read 'Sickling and solubility tests and the peripheral blood film method for screening for sickle cell disease' and not 'Solubility tests and the peripheral blood film method for screening for sickle cell disease'. The title has been corrected on the web version of the article (www.samj.org.za).

In the article entitled 'Non-alcoholic fatty liver disease (NAFLD) in the Western Cape: A descriptive analysis' by F C Kruger *et al.*, which appeared on pp. 168 - 171 of the March 2010 *SAMJ*, the unit for triglycerides in Table I (p. 170) should have been mmol/l and not mg/dl (the figures remain the same). This has been corrected on the web version of the article (www.samj.org.za).

In the 'Guideline on the management of psoriasis in South Africa' (part 2 of the April 2010 *SAMJ*), two errors occurred. On p. 270, in line 5 of the left-hand column, reference 4 should have read 44, and on p. 273, in line 6 of the left-hand column, 2.5 g/d should have read 2.5 mg/d. These have been corrected on the web version of the article (www.samj.org.za).

Withdrawal of published paper

Please note that the article entitled 'Neurocysticercosis in patients presenting with epilepsy at St Elizabeth's Hospital, Lusikisiki' by Ocana GS, Sablon JCO, Tamayo IO, Arena LA, Ocana LMS, Govender S (*S Afr Med J* 2009; 99(8): 588-591) has been withdrawn because key contributors were not included as authors, and material was used without permission.