

CPD questionnaires must be completed online via www.cpdjournals.co.za
After submission you can check the answers and print your certificate.

1. True (A) or false (B):
The leading cause of injury in South Africa is rugby.
2. True (A) or false (B):
The prevalence of fetal alcoholism syndrome in South Africa is the highest in the world.
3. True (A) or false (B):
Women are more likely to be victims of non-sexual violent crime than men.
4. True (A) or false (B):
Young black men are at the greatest risk of homicide.
5. True (A) or false (B):
Schools with clear missions and strong stances against violence have lower rates of bullying.
6. True (A) or false (B):
Within low-income, high-violence neighbourhoods the victims rarely know the perpetrators of violence.
7. True (A) or false (B):
Evidence suggests that violence occurs at higher rates in societies with high levels of economic inequality.
8. True (A) or false (B):
Amitraz is an organic plant food.
9. True (A) or false (B):
Amitraz has alpha-2-adrenergic agonist effects and inhibits monoamine oxidase.
10. True (A) or false (B):
Poisoning by amitraz is always deliberate.
11. Which of the following is FALSE? Patients poisoned by amitraz may present with:
a) Nausea
b) Bradycardia
c) Loss of hair
d) Miosis or mydriasis.
12. True (A) or false (B):
Amitraz and its metabolites or solvents may contribute to a characteristic 'mothball-like' or 'dry-cleaning' odour in the poisoned patient.
13. True (A) or false (B):
The rise of multidrug-resistant bacteria is not of concern because of the rapid production of newer antibiotics.
14. True (A) or false (B):
Appropriate empiric antibiotic guidelines should be based on accurate and up-to-date knowledge of local antibiotic sensitivities.
15. True (A) or false (B):
Unless children attending many South African schools receive food supplementation, malnutrition is likely to impact negatively on their learning ability.
16. True (A) or false (B):
A study showed metal contamination of vegetables grown in a study school garden.
17. True (A) or false (B):
Mortality and morbidity can be reduced in patients presenting with acute myocardial infarction with early interventions such as fibrinolysis.
18. True (A) or false (B):
Maximal benefit of fibrinolysis is seen when the fibrinolytic is given a day after symptom onset.
19. True (A) or false (B):
The study in the Western Cape showed that a key modifiable factor to prolonged door-to-needle times for myocardial infarction was the need for senior review on ECG interpretation.
20. True (A) or false (B):
Patients presenting with atypical symptoms or with undifferentiated chest pain require an ECG to diagnose ST elevation myocardial infarction.

INSTRUCTIONS

1. Read the journal. All the answers will be found there.
2. Go to www.cpdjournals.co.za to answer the questions.

Accreditation number: MDB001/033/10/2011 (Clinical)